

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
6 June 2002 (06.06.2002)

PCT

(10) International Publication Number
WO 02/43647 A2

(51) International Patent Classification⁷: **A61K**

(21) International Application Number: PCT/CA01/01703

(22) International Filing Date:
30 November 2001 (30.11.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/250,131 1 December 2000 (01.12.2000) US
60/327,016 5 October 2001 (05.10.2001) US

(71) Applicant (for all designated States except US): **UNIVERSITY OF OTTAWA** [CA/CA]; 550 Cumberland Avenue, Ottawa, Ontario K1N 6N5 (CA).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BROWN, Earl, Garnet** [CA/CA]; 71 Sunnycrest Drive, Nepean, Ontario K2E 5Y6 (CA). **MBISA, Jean, Lutamyo** [CA/CA]; 1009 Miss Ottawa Street, Gloucester, Ontario K1J 7A9 (CA). **BELL, John, Cameron** [CA/CA]; 248 Knox Crescent, Ottawa, Ontario K1G 0K8 (CA). **STODJL, David, Francis** [CA/CA]; 406-88 Somerset Street, Ottawa, Ontario K2P 0H6 (CA).

(74) Agent: **BERESKIN & PARR**; 40 King Street West, 40th floor, Toronto, Ontario M5H 3Y2 (CA).

(81) Designated States (*national*): AF, AG, AI, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/43647 A2

(54) Title: **ONCOLYTIC VIRUS**

(57) Abstract: Methods of reducing the viability of a tumor cell, infecting a neoplasm in a mammal, utilizing certain non-naturally occurring viruses are disclosed. Viral reassortants, for example reovirus reassortants, and techniques for identifying PKR-sensitive viruses are also disclosed.

ONCOLYTIC VIRUS

SUMMARY OF THE INVENTION

This invention provides methods of reducing the viability of a tumor cell, infecting a neoplasm in a mammal with a virus, or treating a neoplasm in a mammal, comprising administering a non-naturally occurring virus wherein the virus is: a) a reovirus whose mu-2 protein has amino acid residues A, R, M, F, L, M, I, Q, I and S at positions 93, 150, 300, 302, 347, 372, 434, 458, 652 and 726, respectively; or b) a reassortant of two or more parent strains of a viral species selected from the family Reoviridae, or progeny thereof; or c) a virus other than a reovirus wherein the virus other than a reovirus is: i) capable of expressing a reovirus mu-2 protein having amino acid residues A, R, M, F, L, M, I, Q, I and S at positions 93, 150, 300, 302, 347, 372, 434, 458, 652 and 726, respectively, and ii) is a DNA virus, a positive-sense RNA virus, or a negative-sense RNA virus selected from the group consisting of Orthomyxoviridae, Rhabdoviridae and Paramyxoviridae. This invention further provides the use of such non-naturally occurring virus in the manufacture of a medicament for reducing the viability of a tumor cell, infecting a neoplasm in a mammal, or treating a neoplasm in a mammal.

This invention provides a method of identifying a PKR sensitive virus comprising: a) dividing a sample of a virus to be tested into a first portion and second portion; b) contacting PKR +/+ cells with the first portion and contacting PKR -/- cells with the second portion, under conditions permitting growth of the virus in PKR -/- cells; c) determining the rate of growth of the virus in the PKR +/+ cells and in the PKR -/- cells; and d) comparing the growth rates from step c), wherein a higher rate of growth in the PKR -/- cells than in the PKR +/+ cells identifies the virus as PKR sensitive. Such PKR sensitive viruses identified in accordance with this invention are useful for reducing the viability of a tumor cell, infecting a neoplasm in a mammal, or treating a neoplasm in a mammal.

DESCRIPTION OF THE FIGURES

- Figure 1: Virus yield of reovirus strains T1L and T3D in PKR $-/-$ vs. PKR $+/+$ murine embryo fibroblasts.
- Figure 2: Immuno-blot of PKR in MEF Infected with Reo T1L and T3D.
- Figure 3: Lungs of mice with ct26 tumors after treatment with reovirus strains. T1L, T3D, EB96, EB108 and EB146 relative to untreated control lung. The lungs from 2 mice are shown for each treatment.
- Figure 4: The weight of BALB-C mouse lungs relative to the presence of CT26 tumors and reovirus treatment.
- Figure 5: Histological sections stained with hematoxylin and eosin showing lung lobes of mice with ct26 tumors after treatment with reovirus strains. T1L, T3D, EB96, EB108 and EB146 relative to untreated control lung.

DETAILED DESCRIPTION OF THE INVENTION

Throughout this application amino acids are generally identified using the standard one-letter abbreviation, but can also be identified by name or standard three-letter abbreviations.

T3D, T1L, T3A and T2J are standard abbreviations for reovirus strains T3 Dearing, T1 Lang, T3 Abney, and T2 Jones, respectively. The above-listed names of strains and their respective abbreviations are used interchangeably.

As used herein "phenotype" refers to the sequence of the expressed proteins of a virus. In the case of reoviruses the expressed proteins are the gene products of the L1, L2, L3, M1, M2, M3, S1, S2, S3 and S4 genes. Thus, if the amino acid sequences of the products of these genes are the same in two different reoviral strains they are said to have the same phenotype.

As used herein "genotype" refers to the nucleotide sequence of the coding region of a virus. Thus, for example, if the nucleotide sequences of the L1, L2, L3, M1, M2, M3, S1, S2, S3 and S4 genes of two reoviruses are the same in two different reoviral strains they are said to have the same genotype.

The term "PFU" stands for plaque forming units and is a quantitative

measure of live virus particles.

Examples of the anti-neoplastic and anti-tumor methods and use of this invention as described above, include those utilizing a reovirus whose mu-2 protein has amino acid residues A, R, M, F, L, M, I, Q, I and S at positions 93, 150, 300, 302, 347, 372, 434, 458, 652 and 726, respectively. In a more specific embodiment the reoviral mu-2 protein has the amino acid sequence of the mu-2 protein of reovirus strain T3 Dearing, for example when the mu-2 protein is expressed by a gene having the nucleic acid sequence of the M1 gene of reovirus strain T3 Dearing. In a more specific embodiment the reovirus has the same genotype as a reovirus strain selected from the group consisting of eb86, eb129, eb88, eb13, and eb145. In a more specific embodiment the reovirus has a M1 gene whose sequence is the same as the M1 gene of reovirus strain T3 Dearing and an L3 gene whose sequence is the same as the L3 gene of reovirus strain T1 Lang, for example the virus can have the same genotype as a reovirus strain selected from the group consisting of eb28, eb31, eb97, eb123 and g16. In a still more specific embodiment the reovirus has a M1 gene whose sequence is the same as the M1 gene of reovirus strain T3 Dearing and an L3 gene, L1 gene, and S2 gene whose sequences are the same as the corresponding genes of reovirus strain T1 Lang, for example reoviruses having the same genotype as a reovirus strain selected from eb96, eb146 and eb108. In an even more specific embodiment the reovirus has a M1 gene whose sequence is the same as the M1 gene of reovirus strain T3 Dearing and an L3 gene, L1 gene, S2 gene and S4 gene whose sequences are the same as the corresponding genes of reovirus strain T1 Lang, for example reoviruses having the same genotype as reovirus strain eb96.

Other examples of the anti-neoplastic and anti-tumor methods and use of this invention as described above, include those utilizing a virus that is a reassortant of two or more parent strains of a viral species selected from the family Reoviridae, or progeny thereof. For example, reassortants can be made of two, three or four of the reovirus strains T3 Dearing, T1 Lang, T3 Abney, and T2 Jones. In a more specific embodiment the reassortants are generated from parent strains T3 Dearing and T1 Lang. Examples of such strains include eb118, eb73.1, h17, h15, eb39, and h60 as well as the other strains shown in Tables 1 and 2.

Other examples of the anti-neoplastic and anti-tumor methods and use of this invention as described above, include those utilizing a virus other than a reovirus

that is: i) capable of expressing a reovirus mu-2 protein having amino acid residues A, R, M, F, L, M, I, Q, I and S at positions 93, 150, 300, 302, 347, 372, 434, 458, 652 and 726, respectively, and ii) is a DNA virus, a positive-sense RNA virus, or a negative-sense RNA virus selected from the group consisting of the families Orthomyxoviridae, Rhabdoviridae and Paramyxoviridae. Examples of suitable DNA viruses include a Herpesvirus, Adenovirus, Parvovirus, Papovavirus, Iridovirus, Hepadenavirus, Poxvirus, mumps virus, human parainfluenza virus, measles virus or rubella virus. Examples of suitable a positive-sense RNA viruses include a Togavirus, Flavivirus, Picornavirus, or Coronavirus. Examples of suitable negative-sense RNA virus selected from the group consisting of Orthomyxoviridae, Rhabdoviridae and Paramyxoviridae include an influenza virus or a vesicular stomatitis virus.

In accordance with the method of identifying a PKR sensitive virus of this invention as described above, any PKR $+/+$ and $-/-$ cells can be used, and the rate of growth of the virus is determined by any standard technique for monitoring viral growth including those that measure the number of virus particles directly or the quantity of viral proteins. In a specific embodiment the PKR cells are mouse embryo fibroblasts. In another specific embodiment the rate of growth of the virus is determined by a technique selected from the group consisting of plaque titer assay, antibody assay, and Western blot. Each of these techniques is exemplified below. Preferably the growth rate of the virus in PKR $-/-$ cells is at least ten times higher than the growth rate in PKR $+/+$ cells.

In all of the anti-neoplastic and anti-tumor methods and use of this invention as described above, the virus can be a replication competent virus and/or a clonal virus. The virus can be administered by any conventional route, including but not limited to intranasally, intratracheally, intravenously, intraperitoneally or intratumorally. In accordance with the method or use of reducing the viability of a tumor cell described above, the virus can be administered to the tumor cell either *in vivo* or *ex vivo*. When the virus is administered to a mammal, the mammal can be either a human or a non-human mammal such as a mouse, sheep, cow, pig, dog or rabbit. While the optimal dose is expected to differ somewhat from patient to patient and can readily be determined by a skilled clinician, a dosage of from 3×10^7 to 3×10^9 PFU/kg is typical.

The viruses utilized in accordance with this invention can be produced by any conventional means, including reassortment among two or more parent virus strains or the use of standard recombinant genetic techniques. Once produced, such viruses can be reproduced by culturing in cells to produce progeny. The construction of reassortants of viruses is well known and is described, for example in Brown, et al., "The L2 Gene of Reovirus Serotype 3 Controls the Capacity to Interfere, Accumulate Deletions and Establish Persistent Infection" in Double-Stranded RNA Viruses, Compans, et al. eds. Elsevier (1983). For example, reassortants can be made of two, three or four of the reovirus strains T3 Dearing, T1 Lang, T3 Abney, and T2 Jones. Reassortants of T3 Dearing and T1 Lang are described in Example 2. Preferably the virus is replication competent and/or a clonal virus.

This invention will be better understood by reference to the following examples, which illustrate but are not intended to limit the invention described herein.

EXPERIMENTS

Experiment 1: Growth of Reovirus Strains T1L and T3D in PKR Knock-Out and Wild Type Fibroblast Cells

Viral Growth

The effect of PKR on reovirus infection was examined using PKR knock-out (PKR -/-) murine embryo fibroblasts (MEF). Both reovirus T1L and T3D grow to several fold higher titre in PKR -/- relative to PKR +/+ MEF, as measured by plaque assay. (Figure 1) This was associated with a higher percentage of antigen positive cells detected by fluorescent antibody staining described below. Consistent with this, infection of PKR -/- MEF resulted in several fold greater amounts of viral protein as assayed by western blot described below. Although both T1L and T3D grew to higher titres in cells lacking the PKR gene T1L virus grew to higher titres than T3D in either PKR -/- or PKR +/+ cells. (Figure 1)

Indirect Immunostaining

Cells were grown on glass coverslips in 35 mm diameter dishes and were infected with reovirus T1L or T3D at a multiplicity of infection (moi) of 10. After 48 hours incubation the cells were rinsed in PBS and fixed in prechilled acetone for 5 min. After rinsing in PBS (3x5min), 100µl of an appropriate dilution of type-specific rabbit antiviral antisera was applied and incubated at room temperature for 30 min. The coverslips were then rinsed in PBS (3x5min) and treated with the appropriate dilution of Cy3-conjugated donkey anti-rabbit antibody (Jackson ImmunoResearch Laboratories, Inc.) as the secondary antibody. After another 30 min incubation period at room temperature the coverslips were rinsed in PBS (3x5min) and mounted on glass slides in Gel/Mount (Biomedica Corp). All antibody dilutions were done in PBS/3 % BSA.

The samples were examined with a Zeiss microscope equipped with epifluorescence and a 40X 1.40 NA PlanApo objective. The images were collected using Image One Metamorph software and a Hamamatsu chilled charge-coupled digital camera (model C5985). Configuration of the digital images was done using Corel Presentations software.

Immunoblotting

Monolayer cultures of MEF were infected at a moi=10 with T1L or T3D virus as described above. At various times the culture medium was removed and the cells were rinsed with PBS before solubilizing in 1 ml of sample buffer (62.5mM Tris-HCl pH6.8, 10% glycerol, 2 % SDS, 0.05% bromophenol blue and 5 % 2-mercaptoethanol)(Laemmli). Aliquots of 25 µl volume were subjected to SDS PAGE and transblotted onto an Immobilon P membrane (Millipore) at 25V overnight at 4°C. The dried membrane was blocked with 5% (w/v) skim milk powder in PBS for 1hr at RT. This was followed by the addition of type specific rabbit anti-reovirus immune serum as the primary antibody in fresh blocking solution and incubation for 2hr at 4°C. The membrane was then washed three times in PBS and once in TBS (100 mM Tris Hcl pH 7.4, 0.9 % NaCl) to remove phosphate and incubated in 5% milk in TBS containing 1 µg/ml protein A conjugated to alkaline phosphatase obtained from Sigma Chemicals (Oakville, Ont) Finally the membrane was washed 4x in TBS before reaction with chromogenic substrate, nitro blue tetrazolium (NBT) (33 µg/ml) plus 5-bromo-4-chloro-3-indolyl

phosphate (BCIP) (3.3 ul/ml), in alkaline phosphatase buffer (100mM NaCl, 5mM MgCl₂ and 100mM Tris-HCl pH9.5). The reaction was stopped with PBS containing 20mM EDTA.

Experiment 2: Reassortants Between Reovirus Strains T1L and T3D

Production of Genetic reassortants between Reovirus Serotype 1 Lang strain and Serotype 3 Dearing strain.

Mouse L929 cells were coinfectd with Reovirus Serotype 1 Lang strain (T1L) and Serotype 3 Dearing strain (T3D) at a multiplicity of infection of 5 each. Virus was harvested 24 hr post infection by 3 cycles of freezing and thawing before progeny viruses were isolated by 2 cycles of plaque isolation in L929 monolayers. Since each of the corresponding genome segments of T1L and T3D is distinguishable by electrophoretic mobility the genetic composition of each virus was determined by polyacrylamide gel electrophoresis of the segmented double stranded RNA (dsRNA) genome where the mobility of each segment is compared to the parental strains. Gels prepared as described by Laemmli contained 10% polyacrylamide and 0.27% methylene bis-acrylamide. Double-stranded RNA was obtained from L929 cells infected for 3 days and solubilised in buffer containing sodium dodecyl sulphate and was detected in gels stained with ethidium bromide as described previously (Zou S. and E.G. Brown. (1992) Identification of Sequence elements containing signals for replication and encapsidation of the reovirus M1 genome segment. Virology 186:377-88.. The use of this panel of reassortants was first described by E.G. Brown, M. L. Nibert and B.N. Fields (1983) The L2 gene of reovirus serotype 3 controls the capacity to interfere, accumulate deletions and establish persistent infection. in Double-Stranded RNA Viruses. R.W. Compans and D.H.L. Bishop eds. Elsevier Science Publishing Co.

Growth of Reovirus

T1L, T3D and virus stocks from the reassortment procedure described above were prepared in L929 cells grown in Earl's Minimal Essential Medium (MEM) supplemented with 5 % fetal bovine serum and penicillin to 100 units/ml and streptomycin to 100 ug/ml until cytopathic effect was complete. Cells and culture

supernatant were subjected to 3 cycles of freezing and thawing before titration by plaque assay.

Yields in Mouse Embryo Fibroblasts

Wild type PKR^{+/+} cells were obtained from Balb-C mice and PKR^{-/-} cells were obtained from PKR knockout mice. Cell cultures were produced using 15-17 days embryos that had been disaggregated by mincing and trypsin treatment. Cell monolayers were grown in 35 mm plastic dishes in MEM supplemented with 10% FBS and P/S at 37 C in a 5% CO₂ atmosphere. Cells were infected with titrated T1L, T3D or reassortant reovirus at a multiplicity of infection (moi) of 10 by adsorption of stock virus for 0.5 hr with agitation at 15 minute intervals. Unadsorbed virus was removed by 3 washes with 2 ml of warm PBS each before the addition of 3 ml of MEM supplemented with 5 % fetal bovine serum and penicillin to 100 units/ml and streptomycin to 100 ug/ml. The yield of T1L and T3D was assayed at time points over a 4 day period and is shown in Figure 1. Comparison of yields of virus from MEF cells infected with reassortant reovirus was done after 3 days incubation by plaque assay of duplicate cultures. The results are shown below in Table 1 (PKR ^{-/-}) and Table 2 (PKR ^{+/+}).

Plaque assay of reovirus in L929 Cells

Monolayer cultures of L929 cells were decanted of medium and infected in duplicate with 0.1 ml volumes of serially diluted virus in PBS. Virus was adsorbed for 0.5 hr before the application of 3 ml of MEM supplemented with 1 % agar, 5 % FBS and P/S. Cultures were incubated at 37 C and supplementary overlays of 2 ml aliquots of the same medium was added 3 and 6 days post infection. After 8 days of infection the monolayers were stained for 24 hr with 2 ml of the same overlay solution supplemented with neutral red (0.01 % weight/volume) to observe plaques.

Discussion

The genetic basis for the increased ability of T1L to grow in each cell type was determined using T1L x T3D reassortants. The difference in yield in wild type MEF (PKR ^{+/+}) segregated primarily with the M1 gene whereas the difference in yield in PKR ^{-/-} MEF was associated with the L1, L3, M3 and S2 genes and did not

involve the M1 gene. The comparison of the genetic basis for replication in PKR +/+ relative to PKR -/- MEF cells indicates that the ability of the PKR gene to inhibit reovirus infection is dependent on the properties of the M1 gene. Furthermore the extent of replication and thus exploitation of PKR -/- cells is dependent on the nature of the L1, L3, M3 and S2 genes. Thus the reassortant viruses with the greatest differential ability to replicate in PKR -/- relative to PKR +/+ cells possess the T3D M1 gene and the viruses with the greatest ability to replicate in PKR -/- cells (characteristic of many tumor cells) possess the L1, L3, M3 and S2 genes of T1L. Such viruses are restricted in replication of PKR +/+ cells but replicate to a greater extent than either T1L or T3D in PKR -/- cells and are embodied in the properties of the reassortants eb96 and eb108. Statistical analyses of the experimental results are shown in Tables 1, 2 and 3.

The amino acid sequences of the T1L and T3D mu2 proteins are shown in Table 4. Each protein is 736 amino acids long and they differ at 10 aa positions. The observed difference in sensitivity to PKR seen as an ability to replicate in PKR+/+ relative to PKR-/- MEF cells is attributed to the difference in amino acid sequence between these proteins and thus M1 proteins of reoviruses with these amino acid changes or other substitutions at these positions are addressed herein. The mu2 protein is encoded by the M1 gene. The nucleotide sequences of the T1L and T3D M1 gene are shown in Table 5. Each genome segment is 2304 nucleotides long and they differ at 51 nucleotide positions.

TABLE 1: PKR -/-

VIRUS	TITRE	L1	L2	L3	M1	M2	M3	S1	S2	S3	S4	RANK
eb146	7.00E+08	L	L	L	D	L	L	L	L	L	D	1
eb28	5.80E+08	D	D	L	D	D	D	D	L	D	D	2
eb108	4.70E+08	L	D	L	D	L	L	L	L	D	D	3
eb118	4.50E+08	D	D	L	L	D	D	D	D	L	L	4
T1L	4.30E+08	L	L	L	L	L	L	L	L	L	L	5
eb73.1	3.50E+08	L	D	L	L	D	D	D	D	D	D	6
eb31	3.20E+08	L	L	L	D	L	L	L	D	D	L	7
h17	3.00E+08	D	D	L	L	D	D	L	D	D	L	8
H15	2.80E+08	L	D	D	L	D	D	D	D	D	L	9
eb39	2.60E+08	L	D	D	L	D	D	D	D	D	D	10
eb96	1.80E+08	L	D	L	D	L	L	L	L	D	L	11
eb97	1.40E+08	D	D	L	D	D	D	D	D	D	L	12
h60	1.30E+08	D	D	L	L	D	D	D	D	D	L	13
T3D	1.20E+08	D	D	D	D	D	D	D	D	D	D	14
eb123	9.50E+07	D	D	L	D	D	D	D	D	L	D	15
g16	9.30E+07	L	L	L	D	L	L	L	D	L	L	16
eb86	8.50E+07	L	D	D	D	D	L	D	D	D	L	17
eb129	6.30E+07	D	D	D	D	D	L	D	L	L	D	18
eb88	6.00E+07	D	D	D	D	L	D	D	D	D	D	19
eb13	5.30E+07	D	D	D	D	D	D	D	D	D	L	20
eb145	1.30E+07	D	D	D	D	D	L	L	D	D	D	21
t-test		0.045	0.19	0.019	0.024	0.25	0.75	0.57	0.087	0.62	0.76	
M-W test		0.085	0.19	0.007	0.109	0.28	1	0.26	0.047	0.61	0.97	

TABLE 2: PKR +/- (wild type)

VIRUS	TITRE	L1	L2	L3	M1	M2	M3	S1	S2	S3	S4	RANK
h60	3.96E+08	D	D	L	L	D	D	D	D	D	L	1
eb39	2.35E+08	L	D	D	L	D	D	D	D	D	D	2
H15	1.78E+08	L	D	D	L	D	D	D	D	D	L	3
eb118	1.76E+08	D	D	L	L	D	D	D	D	L	L	4
eb146	1.68E+08	L	L	L	D	L	L	L	L	L	D	5
T1L	1.50E+08	L	L	L	L	L	L	L	L	L	L	6
h17	1.46E+08	D	D	L	L	D	D	L	D	D	L	7
eb28	1.30E+08	D	D	L	D	D	D	D	L	D	D	8
eb73.1	1.23E+08	L	D	L	L	D	D	D	D	D	D	9
eb31	5.20E+07	L	L	L	D	L	L	L	D	D	L	10
eb123	4.88E+07	D	D	L	D	D	D	D	D	L	D	11
g16	4.03E+07	L	L	L	D	L	L	L	D	L	L	12
eb129	3.78E+07	D	D	D	D	D	L	D	L	L	D	13
eb97	2.35E+07	D	D	L	D	D	D	D	D	D	L	14
eb96	2.20E+07	L	D	L	D	L	L	L	L	D	L	15
eb108	1.33E+07	L	D	L	D	L	L	L	L	D	D	16
T3D	1.20E+07	D	D	D	D	D	D	D	D	D	D	17
eb13	7.50E+06	D	D	D	D	D	D	D	D	D	L	18
eb86	6.40E+06	L	D	D	D	D	L	D	D	D	L	19
eb88	6.00E+06	D	D	D	D	L	D	D	D	D	D	20
eb145	2.25E+06	D	D	D	D	D	L	L	D	D	D	21
t-test	0.39	0.15	0.056	0.0001	0.68	0.2	0.76	0.56	0.1	0.48		
M-W test	0.4	0.35	0.07	0.0009	0.63	0.21	0.8	0.85	0.24	0.42		

In Tables 1 and 2, parental origin of genome segments is indicated by L (T1L) or D (T3D). Statistical significance was determined using the t-test and the Mann-Whitney (MW) test.

TABLE 3: SUSCEPTIBILITY TO PKR SEGREGATES WITH THE M1 GENE

Gene	Single gene regression (R ² %)		Stepwise regression (R ² %)	
	PKR+/+	PKR-/-	PKR+/+	PKR-/-
L1	0	19 (P=.048)	0	L3 + L1 48 (P=.003)
L3	23.8 (P=.025)	36 (P=.004)	M1+L3 67.0 (P<.001)	36 (P=.004)
M1	51.6 (P<.001)	0	51.6 (P=<.001)	L3 + L1+ M1 56 (P=.0025)
S2	0	16 (P=.073)	0	L3+L1+M3+S 2 63.4 (P<.001)

TABLE 4: Alignment of T1L (GenBank Accession No. CAA42570.1) and T3D (GenBank Accession No. AAA47256.1) mu2 proteins. These amino acid sequences were deduced from cDNA. Each protein is 736 nucleotides long and differs at 10 aa positions.

```

T1L   1
      MAYIAPPAVDSRSSEAIGLLESFGVDAGADANDVSYQDHDYVLDQLQYMLDGYEA
      GDVI 60

Consensus
MAYIAPPAVDSRSSEAIGLLESFGVDAGADANDVSYQDHDYVLDQLQYMLDGYEAGDVI

T3D   1
      MAYIAPPAVDSRSSEAIGLLESFGVDAGADANDVSYQDHDYVLDQLQYMLDGYEA
      GDVI 60


T1L   61
      DALVHKNLHHSVYCLLPKSQLLEYWKSNPSPVDPDNVDRRLRKRLMLKKDLRKDD
      EYNQ 120

Consensus DALVHKNLHHSVYCLLPKSQLLEYWKSNPSP
IPDNVDRRLRKRLMLKKDLRKDEYNQ

T3D   61
      DALVHKNLHHSVYCLLPKSQLLEYWKSNPSPVDPDNVDRRLRKRLMLKKDLRKDD
      EYNQ 120


T1L   121
      LARAFKISDVYAPLISSTTSPMTMIQNLNQGIVYTTTDRVIGARILLYAPRKYA
      STLS 180

Consensus LARAFKISDVYAPLISSTTSPMTMIQNLN
GEIVYTTTDRVIGARILLYAPRKYASTLS

T3D   121
      LARAFKISDVYAPLISSTTSPMTMIQNLNRGEIVYTTTDRVIGARILLYAPRKYA
      STLS 180


T1L   181
      FTMTKCIIPFGKEVGRVPHSRFNVGTFPSIATPKCFVMSGVDIESIPNEFIKLFYQ
      RVKS 240

Consensus
FTMTKCIIPFGKEVGRVPHSRFNVGTFPSIATPKCFVMSGVDIESIPNEFIKLFYQRVKS

T3D   181
      FTMTKCIIPFGKEVGRVPHSRFNVGTFPSIATPKCFVMSGVDIESIPNEFIKLFYQ

```

RVKS 240

T1L 241

VHANILNDISPQIVSDMINRKRLRVHTPSDRRAAQLMHLPHYVKGASHVDVYKVD
VVDV 300

Consensus

VHANILNDISPQIVSDMINRKRLRVHTPSDRRAAQLMHLPHYVKGASHVDVYKVDVD

T3D 241

VHANILNDISPQIVSDMINRKRLRVHTPSDRRAAQLMHLPHYVKGASHVDVYKVD
VVDM 300

T1L 301

LLEVVDVADGLRNVSRKLTMHTVPVCILEMLGIEIADYCIQEDGMFTDWFLLLTM
LSDG 360

Consensus L EVVDVADGLRNVSRKLTMHTVPVCILEMLGIEIADYCIQEDGM
TDWFLLLTMLSDG

T3D 301

LFEVVDVADGLRNVSRKLTMHTVPVCILEMLGIEIADYCIQEDGMLTDWFLLLTM
LSDG 360

T1L 361

LTDRRTHCQYLINPSSVPPDVILNISITGFINRHTIDVMPDIYDFVKPIGAVLPK
GSFKS 420

consensus LTDRRTHCQYL

NPSSVPPDVILNISITGFINRHTIDVMPDIYDFVKPIGAVLPKGSFKS

T3D 361

LTDRRTHCQYLMNPSSVPPDVILNISITGFINRHTIDVMPDIYDFVKPIGAVLPK
GSFKS 420

T1L 421

TIMRVLDSISILGVQIMPRAHVVDSDVEGEQMEPTFEHAVMEIYKGIAGVDSLDDL
IKWV 480

Consensus TIMRVLDSISILG QIMPRAHVVDSDVEGEQMEPTFE
AVMEIYKGIAGVDSLDDLKQV

T3D 421

TIMRVLDSISILGIQIMPRAHVVDSDVEGEQMEPTFEQAVMEIYKGIAGVDSLDDL
IKWV 480

T1L 481

LNSDLIPHDDRIGQLFQAFLPLAKDLLAPMARKFYDNSMSEGRLLTFAHADSELLN
ANYF 540

Consensus

LNSDLIPHDDRLGQLFQAFLPLAKDLLAPMARKFYDNSMSEGRLLTFAHADSELLNANYF

T3D 481

LNSDLIPHDDRLGQLFQAFLPLAKDLLAPMARKFYDNSMSEGRLLTFAHADSELLN
ANYF 540

T1L 541

GHLLRLKIPYITEVNLIRKNREGGELFQLVLSYLYKMYATSAQPKWFGSLLRLLI
CPWL 600

Consensus

GHLLRLKIPYITEVNLIRKNREGGELFQLVLSYLYKMYATSAQPKWFGSLLRLLICPWL

T3D 541

GHLLRLKIPYITEVNLIRKNREGGELFQLVLSYLYKMYATSAQPKWFGSLLRLL
ICPWL 600

T1L 601

HMEKLIGEADPASTSAEIGWHIPREQLMQDGWCGCEDGFIPYVSIRAPRLVMEELM
EKNW 660consensus HMEKLIGEADPASTSAEIGWHIPREQLMQDGWCGCEDGFIPYVSIRAPRLV
EELMEKNW

T3D 601

HMEKLIGEADPASTSAEIGWHIPREQLMQDGWCGCEDGFIPYVSIRAPRLVIEELM
EKNW 660

T1L 661

GOYHAQVIVTDQLVVGEPRRVSAKAVIKGNHLPVKLVSRFACFTLTAKYEMRLSCG
HSTG 720

Consensus

GOYHAQVIVTDQLVVGEPRRVSAKAVIKGNHLPVKLVSRFACFTLTAKYEMRLSCGHSTG

T3D 661

GOYHAQVIVTDQLVVGEPRRVSAKAVIKGNHLPVKLVSRFACFTLTAKYEMRLSCG
HSTG 720

T1L 721 RGAAYNARLAFRSDLA 736

Consensus RGAAY ARLAFRSDLA

T3D 721 RGAAYSARLAFRSDLA 736

TABLE 5: Alignment of the nucleotide sequences of the T1L (GenBank Accession No. X59945.1) and T3D (GenBank Accession No M27261.1) M1 cDNA encoding mu-2 protein. The complete coding sequences are shown. Since reoviruses are double-stranded RNA viruses, the reoviral genome would contain "u" in place to "t". Each genome segment shown below is 2304 nucleotides long that differ at 51 nucleotide positions.

```

T1L   1
      gctattcgcggtcatggcttacatcgagttcctgcggtggtggattcacgttcaa
      gtga 60
      ||||||||||||||||||||||||||||||||||||||||||||||||||||
|||||
T3D   1
      gctattcgcggtcatggcttacatcgagttcctgcggtggtggattcacgttcaa
      gtga 60

T1L 61
      ggctattggactgctagaatcggttggagtagacgctggggctgatgcgaatgacg
      ttcc 120
      ||||||||||||||||||||||||||||||||||||||||||||||||||||
||||||||||||||
T3D 61
      ggctattggactgctagaatcggttggagtagacgctggggctgatgcgaatgacg
      ttcc 120

T1L 121
      atatcaagatcatgactatgtggttgatcagttacagtatatggttagatggatatg
      aggc 180
      ||||||||||||||||||||||||||||||||||||||||||||||||||||
||||||||||||||
T3D 121
      atatcaagatcatgactatgtggttgatcagttacagtatatggttagatggatatg
      aggc 180

T1L 181
      tggcgacgttatcgatgcactcggtccacaagaattggttacatcactccgtctatt
      gctt 240
      ||| ||||||||||||||||||||||||||||||||||||||||||||||||||||
||||||||||||||
T3D 181

```

tggtgacgttatcgatgcactcggtccacaagaattggttacatcactctgtctatt
gctt 240

T1L 241

gttgccacccaaaagtcaactactagagtattggaaaagtaatccttcagtgatac
cgga 300

|||||

|||||

T3D 241

gttgccacccaaaagtcaactattagagtattggaaaagtaatccttcagcgatac
cgga 300

T1L 301

caacggtgatcgtcggcttcgtaaacgactaatgctaaagaaagatctcagaaaag
atga 360

|||||

|||||

T3D 301

caacggtgatcgtcggcttcgtaaacgactaatgctaaagaaagatctcaggaaag
atga 360

T1L 361

tgaataacaatcaactagcgcgtgctttcaagatatcggatgtctacgcacctctca
tctc 420

|||||

|||||

T3D 361

tgaataacaatcagctagcgcgtgctttcaagatatcggatgtctacgcacctctca
tctc 420

T1L 421

atccacgacgtcacccgatgacaatgatccagaacttgaatcaaggcgagatcgtgt
acac 480

|||||

|||||

T3D 421

atccacgacgtcacccgatgacaatgatacagaacttgaatcgaggcgagatcgtgt
acac 480

T1L 481

cacgacggacagggtaattggggctagaatcttggtatatgctcctagaaagtact
atgc 540

|||||

|||||

T3D 481

cacgacggacagggtaattggggctagaatcttggtatatgctcctagaaagtact
atgc 540

T1L 541

gtcaactctatcatttactatgactaagtgcattccgtttggcaaagaggtgg
gtcg 600

|||||

|||||

T3D 541

gtcaactctgtcatttactatgactaagtgcattccgtttggtaaagaggtgg
gtcg 600

T1L 601

tggtcctcactctagatttaatggtggcacatttccatcaattgctaccccgaaat
gttt 660

|||||

|||||

T3D 601

tggttcctcactctcgatttaatggttgccacatttccgtcaattgctaccccgaaat
gttt 660

T1L 661

tgtcatgagtgggggtgatattgagtccatcccaaataaattcatcaagttgtttt
acca 720

|||||

|||||

T3D 661

tgtcatgagtgggggtgatattgagtccatcccaaataaatttatcaagttgtttt
acca 720

T1L 721

gcgcgtcaagagtgttcacgccaatataactaaatgacatatcacctcagatcgtct
ctga 780

|||||

|||||

T3D 721

gcgcgtcaagagtgttcacgctaactataactaaatgacatatctcctcagatcgtct
ctga 780

T1L 781

catgataaacagaaagcgtttgcgcggttcatactccatcagatcgtcgagccgcgc
agtt 840

|||||

|||||

T3D 781

catgataaacagaaagcgtctgcgcggttcatactccatcagatcgtcgagccgcgc
agtt 840

T1L 841

gatgcatttgccttaccatgttaaacgaggagcgtctcacgtcgacgtttacaagg
tgga 900

|||||

|||||

T3D 841

gatgcatttgccttaccatgttaaacgaggagcgtctcacgtcgacgtttacaagg
tgga 900

T1L 901

tggtgtagacgtgttgtagaggtagtggatgtggccgatgggttgcgcaacgta
tctag 960
|||||
|||||

T3D 901

tggtgtagacatgttgttcgaggtagtggatgtggccgatgggttgcgcaacgta
tctag 960

T1L 961

gaaactaactatgcataccgttcggtagtattcttgaaatgttgggtattgaga
ttgc 1020
|||||
|||||

T3D 961

gaaactaactatgcataccgttcctgtatgtattcttgaaatgttgggtattgaga
ttgc 1020

T1L 1021

ggactattgcattcgtcaagaggatggaatgttcacagattggttcctacttttaa
ccat 1080
|||||
|||||

T3D 1021

ggactattgcattcgtcaagaggatggaatgttcacagattggttcctacttttaa
ccat 1080

T1L 1081

gctatctgatggcttaactgatagaaggacgcattgtcaatacttgattaatccgt
caag 1140
|||||
|||||

T3D 1081

gctatctgatggcttgactgatagaaggacgcattgtcaatacttgatgaatccgt
caag 1140

T1L 1141

tgtgcctcctgatgtgatacttaacatctcaattactggatttataaataggcata
caat 1200
|||||
|||||

T3D 1141

tgtgcctcctgatgtgataacttaacatctcaattactggatttataaatagacata
caat 1200

T1L 1201

cgatgtcatgcctgatatatatgacttcgttaaaccattggcgctgtgctgccta
aggg 1260

||||||||||||||

||||||||||||||||||||||||||||||||||||||||

T3D 1201

cgatgtcatgcctgacatatatgacttcgttaaaccattggcgctgtgctgccta
aggg 1260

atcatttaaatcaacaattatgagagttcttgattcaatatcaatattaggaggtcc
agat 1320

.....

1111

atcatttaaatacaacaattatgagagttcttgattcaatatcaatattaggaatcc
aaat 1320

catgccgcgcgccatgtagttagctcggatgaggtgggaggagcaaattggagccta
cgtt 1380

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1

catgccgcgcgccatgtagttgactcagatgaggctgggagcaaatggagccta
cggtt 1380

tgagcatgcggttatggagatatacaaagggattgctggcggttgactcgctggatg
atct 1440

.....

tgagcaggcggttatggagatatacaaagggattgctggcggttgactcgctggatg
atct 1440

catcaagtgggtgctgaactcggatctcattccgcatgatgacaggcttggccaat
tatt 1500

|||||

catcaagtgggtgttgaactcggatctcattccgcatgatgacaggcttggtcaat
tatt 1500

tcaagcggtttctgcctctcgcaaaggacttgtagctccaatggccagaaagtttt
atga 1560

|||||

|||||

T3D 1501

tcaagcggttttgcctctcgcaaaggacttattagctccaatggccagaaagtttt
atga 1560

T1L 1561

taactcaatgagtgagggtagattgctgacattcgctcatgccgacagtgagttgc
tgaa 1620

|||||

|||||

T3D 1561

taactcaatgagtgagggtagattgctaacattcgctcatgccgacagtgagttgc
tgaa 1620

T1L 1621

cgcaaattactttggtcatttattgcgactaaaaataccatatattacagaggtta
atct 1680

|||||

|||||

T3D 1621

cgcaaattactttggtcatttattgcgactaaaaataccatatattacagaggtta
atct 1680

T1L 1681

gatgattcgcaagaatcgtgaggggtggagagctatttcagcttggttatcgtatc
tata 1740

|||||

|||||

T3D 1681

gatgattcgcaagaatcgtgaggggtggagagctatttcagcttggttatcgtatc
tata 1740

T1L 1741

taaaatgtatgctactagcgcgcagcctaataatgggttgatcattattgcgattgt
taat 1800

|||||

T3D 1741

taaaatgtatgctactagcgcgcagcctaataatgggttgatcattattgcgattgt
taat 1800

T1L 1801

atgtccctgggttacatatggagaaattaataggagaagcagacccggcatctacgt
cggc 1860

|||||

T3D 1801

atgtccctgggttacatatggagaaattaataggagaagcagacccggcatctacgt
cggc 1860

T1L 1861

tgaaattggatggcatatccctcgtgaacagctgatgcaagatggatggtgtggat
gtga 1920

|||||

|||||

T3D 1861

tgaaattgggtggcatatccctcgtgaacagctgatgcaagatggatggtgtggat
gtga 1920

T1L 1921

agatggattcattccctatggttagcatacgtgcgccaagactgggttatggaggagt
tgat 1980

||| |||||

|||||

T3D 1921

agacggattcattccctatggttagcatacgtgcgccaagactgggttatagaggagt
tgat 1980

T1L 1981

ggagaagaactggggccaatatcatgcccaagttattgtcactgatcagcttgtcg
tagg 2040

|||||

T3D 1981

ggagaagaactggggccaatatcatgcccaagttattgtcactgatcagcttgtcg
tagg 2040

T1L 2041

cgaaccgcggagggtatctgcccaaggctgtgatcaagggtaatcacttaccagtta
agtt 2100

|||||

|||||

T3D 2041

cgaaccgcggaggggtatctgctaaggctgtgatcaagggtaccacttaccagtta
agtt 2100

T1L 2101

agtttcacgatttgcattgtttcacattgacggcgaagtatgagatgaggctctcgt
gcgg 2160

||||||||||||||||||||||||||||||||||||||||||||||||||||||||||

||||||||

T3D 2101

agtttcacgatttgcattgtttcacattgacggcgaagtatgagatgaggctttcgt
gcgg 2160

T1L 2161

ccatagcactggacggggggctgcatacaatgcgagactagctttccgatctgact
tggc 2220

|||||

|||||

T3D 2161

ccatagcactggacgtggagctgcatacagtgcgagactagctttccgatctgact
tggc 2220

T1L 2221

gtgatccgtgacatgcgtagtgtagacacctgcccctagggtcaatgggggtagggg
cggg 2280

|||||

|||||

T3D 2221

gtgatccgtgacatgcgtagtgtagacacctgctcctagggtcaatgggggtagggg
cggg 2280

T1L 2281 ctaagactacgtacgcgcttcac 2304

|||||

T3D 2281 ctaagactacgtacgcgcttcac 2304

Experiment 3: Assessment of lethal infection in PKR -/- vs. PKR +/+ Mice

Adult Balb-C PKR +/+ or PKR -/- mice were infected with various dosages of infectious reovirus T1L or T3D via the intraperitoneal (IP) or intranasal (IN) route. IP injections involved the administration of 0.1 ml of stock virus or virus diluted in PBS. IN infection involved the application of 0.05 ml volumes of stock virus or virus diluted in PBS onto the nose-pad of mice anaesthetized with halothane (administered at 3% in oxygen). The survival of adult mice was monitored over a 30 day period. Adult PKR +/+ and PKR -/- mice resisted infection with 5×10^6 infectious T3D virus whereas T1L virus killed PKR -/- mice but not PKR +/+ mice at this dose. This demonstrates an enhanced ability of T1L to infect the tissues of PKR -/- mice. Table 5.

Two day old suckling Balb-C PKR +/+ or PKR -/- mice were infected with various dosages of infectious reovirus T1L or T3D via the IN route. IN infectious involved the application of 0.01 ml volumes of stock virus or virus diluted in PBS onto the nose-pad of mice anaesthetized with halothane (administered at 3% in oxygen). The survival of suckling mice was monitored over an 18 day period. Suckling PKR +/+ or PKR -/- mice were both susceptible to similar dosages of T1L whereas T3D virus killed PKR -/- mice much more effectively than PKR +/+ mice, killing them at doses more than 100 fold less than those required to kill wild type suckling mice. This demonstrates an enhanced ability of T3D to infect the tissues of PKR -/- tissues of suckling mice and indicates a difference in the properties of the T1L and the T3D strains with respect to differential replication in PKR +/+ versus PKR -/- mice although both viruses were more restricted in replication of PKR +/+ mice of different ages (adult versus suckling). Table 5.

TABLE 5

ADULT MICE	T1L virus (S/So)		T3D virus (S/So)	
	PKR+/+	PKR-/-	PKR+/+	PKR-/-
5 E6 IP	ND	100 % (3/3)	ND	100 % (3/3)
5 E6 IN	100 % (3/3)	0 % (0/3)	100 % (3/3)	100 % (3/3)
5 E5 IN	ND	100 % (3/3)	ND	ND
SUCKLING MICE				
3 E6 IN	33 % (2/6)	66 % (2/3)	84 % (5/6)	0 % (0/2)
3 E4 IN	100 % (7/7)	ND	100 % (7/7)	0 % (0/4)
3 E3 IN	ND	ND	ND	100 % (3/3)

Experiment 4: Reovirus T3D is a stronger inducer of PKR MEF than T1L

Infection of PKR+/+ MEF results in a greater expression of the phosphorylated form of PKR (Fig. 2). PKR+/+ MEF were infected at a moi of 10 and incubated over a 48 hr period for immunoblot analysis using rabbit anti-PKR serum that reacts with the first 100 amino acids of PKR. Proteins were separated on a 10% polyacrylamide gel and transferred to IMMOBILON membrane (Millipore Inc.) before incubation with 1/100 diluted primary antibody in the presence of casein. After repeated washing the blot was incubated with goat anti-rabbit antibody conjugated with alkaline phosphatase (1/30,000 dilution) (Sigma Inc) for 1 hour before repeated washing and reaction with Attophos substrate for phosphorescent detection as shown in Figure 2. Activation of PKR results in an electrophoretic form of slightly slower mobility indicated as PKR-P. Infection with T3D results in a greater production of this form than with infection with T1L. This demonstrates that PKR expression is enhanced in T3D infected cells and indicates

that this may be responsible for the greater sensitivity of this virus to the PKR gene.

Experiment 5: Proof of principle for Improved Oncolysis of reovirus T1L x T3D
Reassortants: Demonstration that reovirus reassortants with the M1 gene of T3D
and the remaining genes from T1L and T3D have superior oncolytic properties.

Three reassortants were chosen for testing of oncolytic properties relative to their parental viruses. Each of the reassortants, EB96, EB108 and EB146 possessed the M1 gene of T3D and were expected to preferentially replicate in cells that were damaged in their interferon response. These reassortants also possessed their L1, L3 and S2 genes of T1L that would be predicted to provide optimal replication abilities.

Oncolytic testing was performed by intranasal infection of 10^7 pfu of each virus into mice that possessed lung tumors derived from the CT26 colon tumor cell line of Balb-C origin. Adult female BALB-C mice, 4-6 weeks old, were injected in the tail vein with 3×10^5 CT 26 on day 0 of the experiment. On day 7 groups of 3 mice were anaesthetized and infected with 10^7 pfu of virus in a 0.050 volume of culture medium. Mice were housed for an additional 6 days before euthanization with 90% CO₂/10% O₂. Lungs were removed, weighed, fixed in formalin and photographed. One set of lungs was examined histopathologically by hematoxylin and eosin staining after paraffin embedding and sectioning.

The gross appearance of lungs after treatment showed that the untreated control lungs were heavily tumor laden having a pebbled surface appearance due to contiguous tumor nodules (Fig 3). These animals were in the terminal stages of cancer since one animal died at this time and the others were in respiratory distress. These lungs were 3 times heavier than uninfected balb-c lungs indicating the increased tumor mass approximated twice the mass of the lung tissue (Fig 4). Histologically these lungs were covered with a contiguous layer of tumor nodules and internal tumor masses seen as eosinophilic growths of cells (Fig 4 and 5). Infection with T1L virus resulted in a partial freeing of surface tumor growth observable on gross inspection that was also associated with a decrease in interior and surface nodules and a 20 % reduction in lung weight relative to untreated control (Fig 3, 4 and 5). T3D treatment was not as effective as T1L resulting in lungs that were only distinguishable from untreated controls by a slight (8 %)

decrease in size but were similar in gross and microscopic appearance of tumors (Fig 3, 4 and 5).

In dramatic contrast the EB96 reassortant virus cleared the lung of gross tumor mass on treatment (Fig 3). The lungs were of approximately normal weight having been freed of tumor masses (Fig 4). A small number of residual tumor cells remained at this time as detected by histological examination (Fig 5). The lungs were of normal size and appearance except for some circular patterns and dents on the lungs surface that presumably marked the location of prior tumor nodules. EB146 virus was not more effective at tumor lysis than the T3D parental virus (Fig 3, 4 and 5). Reassortant EB108 was partially effective at oncolysis producing results that were marginally better but similar than the T1L parental strain. On comparison of the genotypes of the reassortants it can be seen that the 3 reassortants possess 7 genome segments in common and thus differ in their L2, S3 and S4 genome segments indicating that the latter group of genes include important modulators of oncolysis. The EB96 reassortant is more effective than EB108 solely due to the nature of the S4 gene since these viruses only differ in the parental origin of this gene. This indicates that the T1L S4 gene conferred enhanced oncolytic properties relative to the T3D S4 gene. Since the S4 gene encodes the dsRNA binding protein that blocks PKR activation it is possible that the T1L S4 gene differs in this ability and thus, in concert with other combinations of T1L and T3D genome segments, controls oncolytic potential. In conclusion, the dramatic increase in effectiveness of the EB96 reassortant at oncolysis, relative to the parental T1L and T3D viruses demonstrates the proof of principle that reassortants of reovirus with specific genotypes have enhanced and effective tumor lysis abilities in metastatic tumors in hosts with active immune responses. Table 6.

Table 6: Ranking of the ability of reovirus reassortants to lyse ct26 lung tumors. The relative weight of ct26 tumor bearing lungs relative to untreated control tumor bearing lungs are shown. The parental origin of genome segments are indicated as L for T1L and D for T3D.

VIRUS	TUMOR %	L1	L2	L3	M1	M2	M3	S1	S2	S3	S4	RANK
eb96	41	L	D	L	D	L	L	L	L	D	L	1
eb108	75	L	D	L	D	L	L	L	L	D	D	2
T1L	80	L	L	L	L	L	L	L	L	L	L	3
eb146	89	L	L	L	D	L	L	L	L	L	D	4
T3D	92	D	D	D	D	D	D	D	D	D	D	5

Experiment 6: Ability of T1L x T3D Reassortants to lyse tumors in vitro

A panel of tumor cell lines obtained from the NCI tumor panel (SF539, cns; SKMEL28, melanoma; HT29; NCI H23, nsc-lung; SW620, colon; DU145, prostate) were infected with the T1L, T3D, or the reassortants, EB96, EB108 and EB146 at an moi of 10 and were observed for cytopathic effect over a 5 day period. The ability to lyse tumor cells was scored visually on a scale of – to +++, where – indicates no difference from mock infected cells and +, ++, and +++ indicate 33 % cell destruction, 66 % cell destruction and complete lysis respectively. Although different tumor cell types differed in their susceptibility to lysis by different reovirus parents or reassortants the reassortants viruses were all as effective or more effective than the T3D parental virus at tumor cell lysis in vitro (Table 7).

Table 7: Cytopathology of reovirus T1L and T3D and reassortants in different tumor cell lines

	Tumor cell line					
virus	SF539 Cns	SKMEL28 melanoma	HT29 -	NCI H23 nsc-lung	SW620 colon	DU145 prostate
T1L	++	+++	++	+++	-	++
T3D	-	+++	+	++	-	+
EB96	++	+++	++	++	+	+
EB108	++	+++	++	++	+	+
EB146	++	+++	++	+++	+	++
RAS						

CLAIMS

What is claimed is:

1. A method of reducing the viability of a tumor cell, comprising administering to the tumor cell a non-naturally occurring virus wherein the virus is:
 - a) a reovirus whose mu-2 protein has amino acid residues A, R, M, F, L, M, I, Q, I and S at positions 93, 150, 300, 302, 347, 372, 434, 458, 652 and 726, respectively; or
 - b) a reassortant of two or more parent strains of a viral species selected from the family Reoviridae, or progeny thereof; or
 - c) a virus other than a reovirus capable of expressing a reovirus mu-2 protein having amino acid residues A, R, M, F, L, M, I, Q, I and S at positions 93, 150, 300, 302, 347, 372, 434, 458, 652 and 726, respectively, wherein the virus other than a reovirus is a DNA virus, a positive-sense RNA virus, or a negative-sense RNA virus selected from the group consisting of Orthomyxoviridae, Rhabdoviridae and Paramyxoviridae.
2. A method of infecting a neoplasm in a mammal with a virus, comprising administering to the mammal a non-naturally virus wherein the virus is:
 - a) a reovirus whose mu-2 protein has amino acid residues A, R, M, F, L, M, I, Q, I and S at positions 93, 150, 300, 302, 347, 372, 434, 458, 652 and 726, respectively; or
 - b) a reassortant of two or more parent strains of a viral species selected from the family Reoviridae, or progeny thereof; or

- c) a virus other than a reovirus wherein the virus other than a reovirus is:
 - i) capable of expressing a reovirus mu-2 protein having amino acid residues A, R, M, F, L, M, I, Q, I and S at positions 93, 150, 300, 302, 347, 372, 434, 458, 652 and 726, respectively, and
 - ii) is a DNA virus, a positive-sense RNA virus, or a negative-sense RNA virus selected from the group consisting of Orthomyxoviridae, Rhabdoviridae and Paramyxoviridae.
3. A method of treating a neoplasm in a mammal comprising administering to the mammal a therapeutically effective amount of a non-naturally occurring virus wherein the virus is:
- a) a reovirus whose mu-2 protein has amino acid residues A, R, M, F, L, M, I, Q, I and S at positions 93, 150, 300, 302, 347, 372, 434, 458, 652 and 726, respectively; or
 - b) a reassortant of two or more parent strains of a viral species selected from the family Reoviridae, or progeny thereof; or
 - c) a virus other than a reovirus wherein the virus other than a reovirus is:
 - i) capable of expressing a reovirus mu-2 protein having amino acid residues A, R, M, F, L, M, I, Q, I and S at positions 93, 150, 300, 302, 347, 372, 434, 458, 652 and 726, respectively, and

- ii) is a DNA virus, a positive-sense RNA virus, or a negative-sense RNA virus selected from the group consisting of Orthomyxoviridae, Rhabdoviridae and Paramyxoviridae.
- 4. Use of a non-naturally occurring virus in the manufacture of a medicament for reducing the viability of a tumor cell, infecting a neoplasm in a mammal, or treating a neoplasm in a mammal, wherein the virus is:
 - a) a reovirus whose mu-2 protein has amino acid residues A, R, M, F, L, M, I, Q, I and S at positions 93, 150, 300, 302, 347, 372, 434, 458, 652 and 726, respectively; or
 - b) a reassortant of two or more parent strains of a viral species selected from the family Reoviridae, or progeny thereof; or
 - c) a virus other than a reovirus wherein the virus other than a reovirus is:
 - i) capable of expressing a reovirus mu-2 protein having amino acid residues A, R, M, F, L, M, I, Q, I and S at positions 93, 150, 300, 302, 347, 372, 434, 458, 652 and 726, respectively, and
 - ii) is a DNA virus, a positive-sense RNA virus, or a negative-sense RNA virus selected from the group consisting of Orthomyxoviridae, Rhabdoviridae and Paramyxoviridae.
- 5. The method of claim 1, 2 or 3, or the use of claim 4, wherein the virus is a reovirus whose mu-2 protein has amino acid residues A, R, M, F, L, M, I, Q, I and S at positions 93, 150, 300, 302, 347, 372, 434, 458, 652 and 726, respectively.

6. The method or use of claim 5, wherein the mu-2 protein has the amino acid sequence of the mu-2 protein of reovirus strain T3 Dearing.
7. The method or use of claim 6, wherein the mu-2 protein is expressed by a gene having the nucleic acid sequence of the M1 gene of reovirus strain T3 Dearing.
8. The method of claim 7, wherein the reovirus has the same genotype as a reovirus strain selected from the group consisting of eb86, eb129, eb88, eb13, and eb145.
9. The method or use of claim 7, wherein the reovirus has a L3 gene whose sequence is the same as the L3 gene of reovirus strain T1 Lang.
10. The method or use of claim 9, wherein the reovirus has the same genotype as a reovirus strain selected from the group consisting of eb28, eb31, eb97, eb123 and g16.
11. The method of claim 9, wherein the reovirus has a L1 gene and a S2 gene whose sequences are the same as the corresponding genes of reovirus strain T1 Lang.
12. The method of claim 11, wherein the reovirus has the same genotype as a reovirus strain selected from eb146 and eb108.
13. The method of claim 11, wherein the reovirus has a S4 gene whose sequence is the same as the corresponding gene of reovirus strain T1 Lang.
14. The method of claim 12, wherein the reovirus has the same genotype as reovirus strain eb96.
15. The method of claim 1, 2 or 3 or the use of claim 4, wherein the virus is a reassortant of two or more parent strains of a viral species selected from the

family Reoviridae, or progeny thereof.

16. The method or use of claim 15, wherein the viral species is reovirus and the parent strains are selected from the group consisting of T3 Dearing, T1 Lang, T3 Abney, and T2 Jones.
17. The method or use of claim 16, wherein the parent strains are T3 Dearing and T1 Lang.
18. The method or use of claim 17, wherein the virus is selected from the group consisting of viral strains eb118, eb73.1, h17, h15, eb39, and h60.
19. The method of claim 1, 2 or 3 or the use of claim 4, wherein the virus is a virus other than a reovirus wherein the virus other than a reovirus is:
 - i) capable of expressing a reovirus mu-2 protein having amino acid residues A, R, M, F, L, M, I, Q, I and S at positions 93, 150, 300, 302, 347, 372, 434, 458, 652 and 726, respectively, and
 - ii) is a DNA virus, a positive-sense RNA virus, or a negative-sense RNA virus selected from the group consisting of Orthomyxoviridae, Rhabdoviridae and Paramyxoviridae.
20. The method or use of claim 19, wherein the virus is a DNA virus selected from a Herpesvirus, Adenovirus, Parvovirus, Papovavirus, Iridovirus, Hepadenavirus, Poxvirus, mumps virus, human parainfluenza virus, measles virus or rubella virus.
21. The method or use of claim 19, wherein the virus is a positive-sense RNA virus selected from a Togavirus, Flavivirus, Picornavirus, or Coronavirus.
22. The method or use of claim 19, wherein the virus is a negative-sense RNA virus selected from the group consisting of Orthomyxoviridae,

Rhabdoviridae and Paramyxoviridae.

23. The method or use of claim 19, wherein the virus is an influenza virus or a vesicular stomatitis virus.
24. The method or use of any one of claims 1-23, wherein the virus is a replication competent virus.
25. The method or use of claim 24, wherein the virus is a clonal virus.
26. The method of any one of claims 1-25, wherein the virus is administered by a route selected from the group consisting of intranasally, intratracheally, intravenously, intraperitoneally or intratumorally.
27. The method or use of any one of claims 1-26 wherein the virus is administered to a human or non-human mammal.
28. The method or use of claim 26 or 27 wherein the virus is administered at a dose of from 3×10^7 to 3×10^9 PFU/kg.

1/5




FIGURE 1

2/5




FIGURE 2

3/5





FIGURE 3

4/5

LUNG WEIGHT 6 DAYS POST TREATMENT**FIGURE 4**

5/5




FIGURE 5